



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

STATISTICAL REPORTS OF MORBIDITY AND MORTALITY, STATES AND CITIES OF THE UNITED STATES (untabulated).

FLORIDA.—Week ended April 22, 1911. Reports from the State board of health show diphtheria present in 3 localities with 3 cases, smallpox in 7 counties with 73 cases, malaria in 4 localities with 36 cases, tuberculosis in 8 localities with 12 cases, typhoid fever in 3 localities with 6 cases.

ILLINOIS—*Alton*.—Month of March, 1911. Population, 20,446. Total number of deaths from all causes 32, including tuberculosis 1. Cases reported: Diphtheria 1, scarlet fever 2.

INDIANA—*Indianapolis*.—Two weeks ended April 15, 1911. Population, 233,650. Total number of deaths from all causes 175, including diphtheria 2, scarlet fever 2, tuberculosis 14, typhoid fever 1. Cases reported: Diphtheria 8, measles 61, scarlet fever 27, smallpox 1, tuberculosis 21, typhoid fever 2.

KANSAS.—Month of March, 1911. Population, 1,690,949. Total number of deaths from all causes not reported. Deaths from contagious diseases include diphtheria 6, measles 10, scarlet fever 10, smallpox 2, tuberculosis 64, typhoid fever 4. Cases reported: diphtheria 56, measles 1,314, scarlet fever 363, smallpox 430, tuberculosis 243, typhoid fever 19.

MARYLAND.—Month of February, 1911. Population, 1,295,346. Total number of deaths from all causes, 1,012, including diphtheria 11, measles 3, scarlet fever 4, tuberculosis 119, typhoid fever 13. Cases reported: Diphtheria 65, measles 189, scarlet fever 45, smallpox 4, typhoid fever 56. The typhoid fever cases were distributed as follows: Potomac River watershed 20, Patapsco River watershed 16, Patuxent River watershed 1, Baltimore city water system 1.

NEW YORK—*Rochester*.—Month of March, 1911. Population, 218,149. Total number of deaths from all causes, 314, including diphtheria 4, measles 1, scarlet fever 10, tuberculosis 33, typhoid fever 1. Cases reported: Diphtheria 30, measles 46, scarlet fever 274, tuberculosis, pulmonary, 40, typhoid fever 1.

NORTH DAKOTA.—Month of March, 1911. Population, 577,056. Total number of deaths from all causes, 324, including diphtheria 2. Cases reported: Diphtheria 22, measles 14, scarlet fever 33, smallpox 7, tuberculosis 10, typhoid fever 12.